

sigvaris

MEDICAL COMPRESSION **PRESCRIPTION PAD**

Patient

Date

Diagnosis

Number
of Refills

Compression (mmHg)

15-20 18-25* 20-30 30-40 40-50 50-60

*18-25mmHg only available in Diabetic Compression Sock (Calf only)

Closed Toe Open Toe Patient Choice



Calf
 w/Grip-top



Thigh



Pantyhose



L R
 Thigh w/
waist



Maternity/
Plus Sizes



Wraps
 w/Foot
piece



Armsleeve
 w/Glove
 w/Gauntlet



Separate
Armsleeve Glove



Separate
Armsleeve Gauntlet

SIGVARIS Donning Aid
(see reverse)

Physician
Name

Physician
Signature

Physician
Comments

License
Number

Phone

DISPENSE AS WRITTEN

To find your local SIGVARIS dealer, please visit
sigvaris.com/find or call **1.800.322.7744**

sigvaris

MEDICAL COMPRESSION **PRESCRIPTION PAD**

Donning Accessories

- SIGVARIS ARION Sim-Slide (Open Toe Donning Device)
- SIGVARIS ARION Magnide On/Off (Closed Toe Donning Device)
- Latex-Free Donning Gloves
- SIMON (Donning Device)
- Rubber Donning Gloves
- SIGVARIS Doff N' Donner (Donning Device)
- Cotton Underliner
- S.O.S. (Slip On SIGVARIS Donning Device)

Compression Symptoms / Indications

15–20mmHg	Heavy, fatigued, tired legs · Mild edema · Prophylaxis of varicosities & swelling during pregnancy · Prophylaxis for upper arm edema · Reduce swelling · Prophylaxis of DVT during travel
18–25mmHg	Diabetics who experience edema (see Cautions below)
20–30mmHg	Aching legs · Mild varicosities with minimal edema · Mild varicosities during pregnancy · Mild-moderate edema upper extremities · Post-procedure of small veins · High risk of VTE · Moderate varicosities with moderate edema (also during pregnancy)
30–40mmHg	Post-procedure of larger veins · CEAP C3, C4, C5, C6* · DVT or Post-Thrombotic Syndrome · Superficial phlebitis · Lymphedema after decongestant therapy · Orthostatic /postural hypotension · Post-traumatic edema
40–50mmHg or higher	Severe varicosities · Severe edema · Lymphedema after decongestant therapy · CEAP C4, C5, C6* · Severe post-traumatic edema · Severe Post-Thrombotic Syndrome

Containment Symptoms / Indications

Low Mild to moderate edema

High Moderate to severe edema · Lymphedema · Fibrotic tissue

Contra-indications Arterial insufficiency (ABI <0.5), intermittent claudication, ischemia · Uncontrolled congestive heart failure · Acute dermatitis, weeping dermatosis, cutaneous sepsis · See product insert for full information

Precautions Signs of infection · Skin sensitivities or allergies · Neuropathy · Diabetes with advanced · arterial compromise · Confinement to bed or non-ambulatory use unless otherwise prescribed by the physician

No liability accepted for non-observance of contra-indications and cautions.

* Fronck HS, Bergan JJ, et al. The Fundamentals of Phlebology; Venous Disease for Clinicians. 2004. pg 151

All Sigvaris medical measured graduated compression garments are guaranteed to retain their stated compression for 6 months with proper care.

sigvaris.com

SIGVARIS and SIGVARIS GROUP are registered trademarks of SIGVARIS, CH-9014 St.Gallen/Switzerland, in many countries worldwide. © Copyright 2019 SIGVARIS, Inc.